


**WHAT IS CLAIMED IS:**

1. Chemokine peptide 3, a variant, or a derivative thereof.
2. Chemokine peptide 2, a variant, or a derivative thereof.
3. The peptide of claim 1 wherein the chemokine is not IL8 or NAP-2.
4. The peptide of claim 1 which is a variant of peptide 3[MCP-1].
5. The peptide of claim 4 which is leu<sub>4</sub>ile<sub>11</sub>peptide 3(3-12)[MCP-1].
6. The peptide of claim 1 or 2 which is a CC chemokine.
7. The peptide of claim 6 wherein the CC chemokine is MCP-1, RANTES, MCP-2, MCP-3, MCP-4, eotaxin, MIP1 $\alpha$ , MIP1 $\beta$ , LARC, I309, HCC-1, TARC or Ck $\beta$ 8.
8. The peptide of claim 1 or 2 which is a CXC chemokine.
9. The peptide of claim 8 wherein the CXC chemokine is IP-10, PF-4, SDF-1, NAP-2, GRO $\alpha$ , GRO $\beta$ , GRO $\gamma$  or ENA78.
10. The peptide of claim 8 wherein the CXC chemokine is IL-8, IP-10, SDF-1, PF-4, NAP-2, GRO $\alpha$ , GRO $\beta$ , GRO $\gamma$ , NAP-2 or ENA78.
11. A CRD derivative of chemokine peptide 3 or a variant thereof.

12. The derivative of claim 11 which is CRD-Cys<sub>13</sub>leu<sub>4</sub>ile<sub>11</sub>peptide 3(3-12)[MCP-1].
13. A CRD derivative of chemokine peptide 2 or a variant thereof.
14. An isolated and purified nucleic acid molecule comprising a preselected nucleic acid segment encoding the peptide of claim 1 or 2.
15. An isolated and purified nucleic acid molecule comprising a preselected nucleic acid segment which is the complement of the nucleic acid segment of claim 14.
16. A compound of formula (IV):



wherein R<sup>1</sup> is aryl, heteroaryl, coumaryl or chromanyl; wherein R<sup>2</sup> is N(R<sup>a</sup>)(R<sup>b</sup>); wherein R<sup>3</sup> is N(R<sup>c</sup>)(R<sup>d</sup>); wherein Y is oxo or thioxo; wherein Z is (C<sub>1</sub>-C<sub>10</sub>)alkyl; wherein R<sup>a</sup>-R<sup>d</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>1</sub>-C<sub>10</sub>)alkanoyl, phenyl, benzyl or phenethyl; or wherein R<sup>a</sup> and R<sup>b</sup>, or R<sup>c</sup> and R<sup>d</sup>, together with the nitrogen to which they are attached form a pyrrolidino, piperidino or morpholino ring; or a pharmaceutically acceptable salt thereof.

17. A compound of formula (V):

wherein  $R^4$  is  $NR_kR_l$ ; wherein  $R^5$  is  $NR_mR_n$ ; wherein  $R^6$  is  $NR_oR_p$ ; wherein  $R^7$  is  $NR_qR_r$ ; wherein  $R^8$  is hydrogen, hydroxy,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkyl,  $(C_1-C_{10})$ alkoxy,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkoxy,  $NR_sR_t$ , the N-terminal residue of an amino acid or a peptide of 2 to about 25 amino acid residues; wherein  $R_k$ ,  $R_l$ ,  $R_o$ , and  $R_p$  are each hydrogen; wherein  $R_m$  and  $R_n$  are each independently hydrogen, acetyl,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl, propoxy, butoxy, *tert*-butoxycarbonyl, 9-fluorenylmethoxycarbonyl, the C-terminal residue of an amino acid or a peptide of 2 to about 25 amino acid residues; wherein  $R_q$  and  $R_r$  are each independently hydrogen,  $(C_1-C_{10})$ alkyl, or  $(C_3-C_6)$ cycloalkyl; and wherein  $R_s$  and  $R_t$  are each independently hydrogen,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkyl, phenyl, benzyl, or phenethyl; or a pharmaceutically acceptable salt thereof.

18. A method of preventing or inhibiting an indication associated with a chemokine-induced activity, comprising: administering to a mammal afflicted with, or at risk of, the indication an amount of a chemokine peptide 3, a variant thereof, a derivative thereof, or a combination thereof, effective to prevent or inhibit said activity, wherein the chemokine is not IL8 or NAP-2.

19. A method to inhibit the activity of more than one chemokine, comprising: administering to a mammal in need thereof an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
20. A method to increase or enhance a chemokine-associated inflammatory response in a mammal, comprising: administering to the mammal an amount of a chemokine peptide 2, a variant thereof, a derivative thereof, or a combination thereof effective to increase or enhance said response.
21. A method of preventing or inhibiting an indication associated with monocyte or macrophage recruitment, comprising: administering to a mammal at risk of, or afflicted with, the indication an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
22. A method of preventing or inhibiting an indication associated with histamine release from basophils or mast cells, comprising administering to a mammal at risk of, or afflicted with, the indication an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
23. A method to modulate the chemokine-induced activity of a macrophage at a preselected physiological site, comprising: administering to a mammal a dosage form comprising an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof, wherein the dosage form is linked to a site targeting moiety.


24. A method to augment an immune response, comprising: administering to a mammal an immunogenic moiety and an amount of a chemokine peptide 2, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof, wherein the amount is effective to augment the immune response of the mammal to the immunogenic moiety.
25. A therapeutic method to prevent or treat a vascular indication, comprising: administering to a mammal in need of such therapy an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof, wherein the indication is coronary artery disease, myocardial infarction, unstable angina pectoris, atherosclerosis or vasculitis.
26. A therapeutic method to prevent or inhibit lentiviral infection or replication, comprising: administering to a mammal in need of such therapy an effective amount of a chemokine peptide 2, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
27. The method of claim 26 wherein the lentivirus is HIV.
28. The method of claim 27 further comprising administering an antiviral agent before, during and/ or after the administration of the peptide, a variant thereof, derivative thereof, the compound of formula (IV) or the compound of formula (V).
29. A therapeutic method to prevent or treat low bone mineral density, comprising: administering to a mammal in need of such therapy an effective

amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.

30. A method of inhibiting a parasitic infection in a vertebrate animal, comprising: administering to the animal an effective amount of a chemokine peptide 2, a variant thereof, a derivative thereof, or a combination thereof.
31. The method of claim 30 wherein the animal is a human with malaria.
32. A therapeutic method to prevent or treat an autoimmune disease, comprising: administering to a mammal in need of such therapy an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
33. A method of suppressing tumor growth in a vertebrate animal, comprising: administering to said vertebrate an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
34. A method for preventing or treating psoriasis in a mammal, comprising: administering to the mammal an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
35. A method to increase or enhance macrophage-associated activity at a tumor site, comprising: administering an effective amount of a chemokine peptide


3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.

36. A method to enhance wound healing, comprising: administering an effective amount of a chemokine peptide 3, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
37. A method of treating a mammal afflicted with, or at risk of, an indication associated with chemokine-induced activity, comprising: administering to the mammal an effective amount of a compound of formula (IV):



wherein R<sup>1</sup> is aryl, heteroaryl, coumaryl or chromanyl; wherein R<sup>2</sup> is N(R<sup>a</sup>)(R<sup>b</sup>); wherein R<sup>3</sup> is N(R<sup>c</sup>)(R<sup>d</sup>); wherein Y is oxo or thioxo; wherein Z is (C<sub>1</sub>-C<sub>10</sub>)alkyl; wherein R<sup>a</sup>-R<sup>d</sup> are each independently hydrogen, (C<sub>1</sub>-C<sub>10</sub>)alkyl, (C<sub>1</sub>-C<sub>10</sub>)alkanoyl, phenyl, benzyl or phenethyl; or wherein R<sup>a</sup> and R<sup>b</sup>, or R<sup>c</sup> and R<sup>d</sup>, together with the nitrogen to which they are attached form a pyrrolidino, piperidino or morpholino ring; or a pharmaceutically acceptable salt thereof.

38. A method of treating a mammal afflicted with, or at risk of, an indication associated with chemokine-induced activity, comprising: administering to the mammal an effective amount of a compound of formula (V):



wherein  $R^4$  is  $NR_kR_l$ ; wherein  $R^5$  is  $NR_mR_n$ ; wherein  $R^6$  is  $NR_oR_p$ ; wherein  $R^7$  is  $NR_qR_r$ ; wherein  $R^8$  is hydrogen, hydroxy,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkyl,  $(C_1-C_{10})$ alkoxy,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkoxy,  $NR_sR_t$ , the N-terminal residue of an amino acid or a peptide of 2 to about 25 amino acid residues; wherein  $R_k$ ,  $R_l$ ,  $R_o$ , and  $R_p$  are each hydrogen; wherein  $R_m$  and  $R_n$  are each independently hydrogen, acetyl,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl, propoxy, butoxy, *tert*-butoxycarbonyl, 9-fluorenylmethoxycarbonyl or the C-terminal residue of an amino acid or a peptide of 2 to about 25 amino acid residues; wherein  $R_q$  and  $R_r$  are each independently hydrogen,  $(C_1-C_{10})$ alkyl, or  $(C_3-C_6)$ cycloalkyl; and wherein  $R_s$  and  $R_t$  are each independently hydrogen,  $(C_1-C_{10})$ alkyl,  $(C_3-C_6)$ cycloalkyl,  $(C_3-C_6)$ cycloalkyl $(C_1-C_6)$ alkyl, phenyl, benzyl, or phenethyl; or a pharmaceutically acceptable salt thereof.

39. An immunogenic composition comprising an immunogenic moiety and an amount of a chemokine peptide 2, a variant thereof, a derivative thereof, a compound of formula (IV), a compound of formula (V), or a combination thereof.
40. A method of preventing or inhibiting an indication associated with a chemokine-induced activity, comprising: administering to a mammal



afflicted with, or at risk of, the indication an amount of the nucleic acid molecule of claim 14 effective to prevent or inhibit said activity.

41. A method of preventing or inhibiting an indication associated with a chemokine-induced activity, comprising: administering to a mammal afflicted with, or at risk of, the indication an amount of the nucleic acid molecule of claim 15 effective to prevent or inhibit said activity.

add  
B5

08927939-091197  
26FF60-6E622680